


8000 GENERAL TRAUMA CARE



8000 GENERAL TRAUMA CARE

EMR	EMT	EMT-IV	AEMT	INTERMEDIATE	PARAMEDIC
-----	-----	--------	------	--------------	-----------

FULL Trauma Team Criteria

Persons who sustain Injury with any of the following:

PRIMARY SURVEY:								
	Adult (15+)	Child (0-14)						
Airway	Unable to adequately ventilate. Intubated or assisted ventilation.	Unable to adequately ventilate. Intubated or assisted ventilation.						
Breathing	Resp. rate < 10 or > 30/min.	Any sign of resp. insufficiency (hypoxia, accessory muscle use, grunting)						
Circulation	Systolic Blood Pressure (BP/SBP) < 90 mmHg or < 110 for age ≥ 65	Any sign of abnormal perfusion (cap refill > 2sec; BP low for age)						
		<table border="1"> <thead> <tr> <th>Age</th> <th>SBP (mm/Hg)</th> </tr> </thead> <tbody> <tr> <td>< 1 yr.</td> <td>< 60</td> </tr> <tr> <td>1 – 10 yrs.</td> <td>< 70 + 2x</td> </tr> <tr> <td>> 10 yrs.</td> <td>< 90</td> </tr> </tbody> </table>	Age	SBP (mm/Hg)	< 1 yr.	< 60	1 – 10 yrs.	< 70 + 2x
Age	SBP (mm/Hg)							
< 1 yr.	< 60							
1 – 10 yrs.	< 70 + 2x							
> 10 yrs.	< 90							
Deficit	GCS Motor Score ≤ 5 OR Total GCS Score ≤ 8	AVPU – Resp. to Pain or Unresponsive						
<input type="checkbox"/> Deterioration of previously stable patient that meets criteria <input type="checkbox"/> Transfers requiring blood transfusion to maintain vital signs <input type="checkbox"/> ED physician discretion								

SECONDARY SURVEY: ANATOMIC
Penetrating injuries to the head, neck, torso or extremities proximal to elbow/knee
Open or depressed skull fracture
Paralysis or neurological deficit, suspected spinal cord injury
Flail Chest (chest wall instability or deformity)
Unstable pelvic fracture
Amputation proximal to the wrist or ankle
Two or more proximal long bone fractures (humerus or femur)
Crushed, devalved, mangled, or pulseless extremity



- Full Trauma Team activation for any pt with a tourniquet in place
- Full Trauma Team activation for any pt with a pelvic binder.

EMR	EMT	EMT-IV	AEMT	INTERMEDIATE	PARAMEDIC
-----	-----	--------	------	--------------	-----------

LIMITED Trauma Team Criteria

Persons who sustain injury within 24 hours with any of the following:

MECHANISM OF INJURY

Falls (adult): > 20 ft.; **or** (child): > 15 ft. or 3x height

High risk auto crash, with:

- Intrusion of vehicle \leq 12" in occupant compartment;
or > 18" other site
- Ejection (partial or complete) from automobile
- Death in same passenger compartment

Auto versus pedestrian/cyclist thrown, run over **or** with significant impact



Motorcycle crash > 20 mph

High energy dissipation or rapid decelerating incidents, ie:

- Ejection from motorcycle, ATV, animal, etc.
- Striking fixed object with momentum
- Blast or explosion

High energy electrical injury

Burns > 10% TBSA (2° or 3°) and/or inhalation injury

Suspected non-accidental trauma

EMS provider judgment

Version 10.2020

uchealth